

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 August 2002 (22.08.2002)

PCT

(10) International Publication Number
WO 02/064160 A2

(51) International Patent Classification⁷: **A61K 38/31**,
31/404, A61P 43/00

(US). **DANESI, Romano** [Stateless/IT]; I-Pisa (IT).
BOCCI, Guido [Stateless/IT]; ** (IT). **DELTACCA,**
Mario [Stateless/IT]; ** (IT).

(21) International Application Number: PCT/US02/01125

(74) Agent: **CLARK, Paul, T.**; Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110 (US).

(22) International Filing Date: 14 January 2002 (14.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/261,439 12 January 2001 (12.01.2001) US
60/275,294 13 March 2001 (13.03.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 60/261,439 (CIP)
Filed on 12 January 2001 (12.01.2001)
US 60/275,294 (CIP)
Filed on 13 March 2001 (13.03.2001)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): SOCI-
ETE DE CONSEIL DE RECHERCHES ET D'APPLI-
CATIONS SCIENTIFIQUES (S.C.R.A.S.) [Stateless/-];
** (-).

Published:

— without international search report and to be republished
upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DEWITT**
CULLER, Michael [Stateless/US]; Hopkinton, MA

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHARMACEUTICAL COMPOSITIONS WHICH INHIBIT VASCULAR PROLIFERATION AND METHOD OF USE THEREOF

(57) Abstract: The present invention relates to a method of treating vascular proliferation in a patient in need thereof. The method includes the step of administering a therapeutically effective amount of a type-1 somatostatin agonist to said patient.



WO 02/064160 A2

5

**PHARMACEUTICAL COMPOSITIONS WHICH INHIBIT
VASCULAR PROLIFERATION AND METHOD OF USE THEREOF**

BACKGROUND OF THE INVENTION

Angiogenesis, the development of new capillaries from preexisting blood
10 vessels, is a critical process in the progression of solid neoplasms and in many other
pathological conditions such as diabetic retinopathy and rheumatoid arthritis (Folkman,
J., *Nature Medicine* 1995, 1: 27-31). Different strategies to target the vascular
development have been extensively studied and the availability of reliable *in vitro* model
systems in model systems in angiogenesis research has been crucial for the study of
15 specific inhibitors (Jain, R.K., *et al.*, *Nature Medicine* 1997, 3: 1203-1208).

It is now widely recognized that the ability of a tumor to induce proliferation of
new blood vessels from its host has a profound effect on cancer growth and metastasis.
The process of tumor angiogenesis is mediated by a balance of positive and negative
regulators of microvessel growth and the development of new blood vessels may be
20 divided into three different sequential steps: 1) cell-mediated, proteolytic degradation of
the basement membrane; 2) endothelial cell migration and proliferation out of the vessel
into the surrounding extracellular matrix; 3) organization of the cells into tube-like
structures (Folkman, J., *Nature Medicine* 1995, 1: 27-31).

Somatostatin (somatotropin release inhibiting factor or SRIF) has both a 14
25 amino acid isoform (somatostatin-14) and a 28 amino acid isoform (somatostatin-28).
See Wilson, J. & Foster, D., *Williams Textbook of Endocrinology*, p. 510 (7th ed., 1985).

The compound is an inhibitor of secretion of the growth hormone and was originally
isolated from the hypothalamus. Brazeau, *et al.*, *Science* 179:77 (1973). Native
somatostatin has a very short duration of effect *in vivo* since it is rapidly inactivated by
30 endo- and exopeptidase. Many novel analogs (e.g., peptide and non-peptide
compounds) have been prepared in order to enhance the duration of effect, biological
activity, and selectivity (e.g., for the particular somatostatin receptor) of this hormone.
Such analogs of somatostatin will be called "somatostatin agonists" herein.

Various somatostatin receptors (SSTRs) have been isolated, e.g., SSTR-1,
35 SSTR-2, SSTR-3, SSTR-4, and SSTR-5. Thus, a somatostatin agonist may be a SSTR-
1 agonist, and/or a SSTR-2 agonist, and/or a SSTR-3 agonist, and/or a SSTR-4 agonist
and/or a SSTR-5 agonist.

The antiangiogenic activity of somatostatin analogues has been previously demonstrated in some *in vitro* and *in vivo* experimental models (Danesi, R. *et al.*, Clinical Cancer Research 1997, 3: 265-272; Woltering, E.A., *et al.*, Investigational New Drug 1997, 15: 77-86). In addition to this, long-term octreotide treatment was able to
5 reduce the progression of neovascularization associated with severe proliferative retinopathy in diabetic patients (Mallet *et al.*, 1992).

The determination of which somatostatin subtype or subtypes are involved in the antiangiogenic property of somatostatin would allow for the development of therapeutic compositions with maximum efficacy and minimum side effects. However, previous
10 studies in this field have resulted in contradictory and/or inconclusive findings regarding the role which each of the five somatostatin receptor subtypes may play in respect of the antiangiogenic activity of somatostatin.

SUMMARY OF THE INVENTION

The present invention relates to compositions comprising a somatostatin type-1
15 receptor agonist which are useful for the inhibition of vascular proliferation in a subject. The present invention further relates to a method of treating vascular proliferation, e.g., angiogenesis and restenosis, in a patient (e.g., a mammal such as a human) in need of such treatment. The method includes the step of administering a therapeutically effective amount of a somatostatin type-1 receptor (SSTR-1) agonist (e.g., a
20 somatostatin type-1 selective agonist) to said patient.

The present invention also relates to a method of inhibiting smooth muscle proliferation, endothelial cell proliferation, and new blood vessel sprouting in a patient in need of such inhibition. The method includes the step of administering a therapeutically effective amount of a somatostatin type-1 receptor (SSTR-1) agonist (e.g., a
25 somatostatin type-1 selective agonist) to said patient.

Examples of clinical indications which can be treated by the present invention include, but are not limited to, autoimmune diseases (e.g., arthritis, scleroderma, etc.), cancerous tumors, corneal graft neovascularization, diabetic retinopathy, hemangioma, hypertrophic scars, and psoriasis, as well as vascular proliferation associated with
30 surgical procedures, e.g., angioplasty and AV shunts.

Further examples of disease states that may be amenable to treatment with the subject therapeutic composition and method of the invention are, in respect of the skin: warts, granulomas, Kaposi's sarcoma, allergic oedema, and the like; in respect of the uterus and ovary: endometriosis, dysfunctional uterine bleeding, follicular cysts, and the
35 like; in respect of the eye: retinopathy of prematurity, choroidal and other intraocular disorders, macular degeneration, age-related macular degeneration, and the like.

Further examples of disease states that may be amenable to treatment with the subject therapeutic composition and method of the invention are disclosed in Folkman, J., *Seminars in Medicine of the Beth Israel Hospital*, Boston, Vol. 333, No 26, pp. 1757-1763.

5 Indeed as is well known in the art the list of known diseases and conditions for which an agent capable of inhibiting angiogenesis, smooth muscle proliferation, endothelial cell proliferation, and/or new blood vessel sprouting is quite substantial, and includes, without limitation: solid tumors, tumor metastasis, benign tumors, for example, acoustic neuromas, neurofibromas, and trachomas, leukemia, pyogenic granuloma, 10 myocardial angiogenesis, plaque neovascularization, atherosclerosis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, ocular and corneal angiogenic conditions, for example, corneal graft rejection, Osler-Webber Syndrome, rubeosis, neovascular glaucoma, retrolental fibroplasia, and diabetic retinopathy, diabetic neovascularization, wound healing, 15 fractures, vasculogenesis, hematopoiesis, ovulation, menstruation, placentation, cat scratch disease (*Rochelle minalia quintosa*), ulcers (peptic; *Helicobacter pylori*), psoriasis, (including, e.g., telangiectasia psoriasis), rheumatoid arthritis, Crohn's disease, intestinal adhesions, scarring, (i.e., formation of high density tissue including cells and connective tissue), hypertrophic scars, (i.e., keloids), telangiectasia; 20 hemophiliac joints; angiofibroma; and wound granulation. These diseases and conditions are discussed in detail in the literature, for example, as in the following U.S. patents and International Patent Publications.: 6,323,228, 6,294,532, 6,288,228, 6,288,024, 6,284,726, 6,280,739, 6,265,407, 6,265,403, 6,258,812, 6,255,355, 6,255,353, 6,251,867, 6,242,481, 6,235,756, 6,235,741, 6,228,879, 6,228,871, 25 6,225,340, 6,214,800, 6,201,104, 6,177,401, 6,174,861, 6,150,407, 6,150,362, 6,117,862, 6,114,355, 6,090,794, 6,086,865, 6,071,948, 6,057,290, 6,057,122, 6,028,061, 6,025,353, 6,025,331, 6,024,688, 6,017,949, 5,997,868, 5,994,388, 5,994,292, 5,990,280, 5,985,878, 5,985,330, 5,981,484, 5,972,922, 5,972,896, 5,948,403, 5,932,611, 5,902,790, 5,874,081, 5,847,002, 5,843,925, 5,837,680, 30 5,807,731, 5,801,146, 5,766,591, 5,753,230, 5,744,492, 5,733,876, 5,721,226, 5,712,291, 5,698,586, 5,696,147, 5,677,181, 5,646,136, 5,629,340, 5,629,327, 5,610,166, 5,593,990, 5,574,026, 5,567,693, 5,567,417, 5,563,130, 5,512,550, 5,506,208, WO02/02609, WO02/02593, WO02/00877, WO02/00690, WO02/00017, WO01/93806, WO01/85796, WO01/81579, WO01/81311, WO01/79157, WO01/74299, 35 WO01/72699, WO01/72297, WO01/66127, WO01/62799, WO01/62725, WO01/59100, WO01/58899, WO01/51048, WO01/46110, WO01/45751, WO01/35977, WO01/34195, WO01/29085, WO01/28577, WO01/25433, WO01/23375, WO01/21831, WO01/19987,

WO01/19868, WO01/12809, WO01/12226, WO01/12210, WO01/10859, WO01/09113,
WO01/07057, WO01/04157, WO01/03720, WO01/00201, WO00/75124, WO00/73445,
WO00/73340, WO00/59532, WO00/54770, WO00/54762, WO00/53757, WO00/53753,
WO00/53752, WO00/52158, WO00/48591, WO00/47212, WO00/47193, WO00/43393,
5 WO00/40597, WO00/35407, WO00/32221, WO00/32180, WO00/30628, WO00/27866,
WO00/27415, WO00/27340, WO00/24415, WO00/21561, WO00/20577, WO00/20026,
WO00/19995, WO00/15792, WO00/12089, WO00/10507, WO00/10506, WO00/09657,
WO00/09495, WO00/05356, WO00/02902, WO00/02871, WO00/02585, WO00/01383,
WO99/62549, WO99/61590, WO99/61432, WO99/60984, WO99/58139, WO99/48495,
10 WO99/45909, WO99/37776, WO99/31088, WO99/26622, WO99/26480, WO99/23105,
WO99/22760, WO99/16755, WO99/16465, WO99/14234, WO99/10349, WO99/09982,
WO99/04806, WO99/04803, WO98/58929, WO98/58919, WO98/54093, WO98/51326,
WO98/41205, WO98/36760, WO98/35958, WO98/31688, WO98/19712, WO98/19649,
WO98/17796, WO98/13071, WO98/12226, WO98/05323, WO98/05293, WO97/45137,
15 WO97/41824, WO97/35567, WO97/32583, WO97/30085, and WO97/26258. The
contents of each of the foregoing patents and patent publications is hereby incorporated
by reference in its entirety.

Definitions of "somatostatin type-1 receptor agonist" and "somatostatin type-1
receptor selective agonist" are provided below. A therapeutically effective amount
20 depends upon the condition being treated, the route of administration chosen, and the
specific activity of the compound used and ultimately will be decided by the attending
physician or veterinarian (e.g., between 5 g/day and 5 mg/day). In one embodiment, the
somatostatin agonist is administered to the patient until the condition being treated has
subsided. In another embodiment, the somatostatin agonist is administered for the
25 lifetime of the patient.

The somatostatin agonist may be injected parenterally, e.g., intravenously, into
the bloodstream of the subject being treated. However, it will be readily appreciated by
those skilled in the art that the route, such as intravenous, subcutaneous, intramuscular,
intraperitoneal, enterally, transdermally, transmucously, sustained released polymer
30 compositions (e.g., a lactic acid polymer or lactic acid and glycolic acid copolymer
microparticle or implant), perfusion, nasal, oral, etc., will vary with the condition being
treated and the activity and bioavailability of the somatostatin agonist being used.

The somatostatin agonist may also be provided as a coating or as a component
of a coating on a surface of a device implanted within the body. For example, the
35 provision of the somatostatin agonist within or upon a vascular stent is useful for the
treatment of restenosis which is often associated with stent implantation.

The SSSTR-1 agonist can be administered systemically and/or locally or topically, as needed. For prevention of adhesions, the SSSTR-1 agonist would typically be applied at the time of surgery, preferably in a controlled release formulation and/or using barrier technology.

5 While it is possible for the somatostatin agonist to be administered as the pure or substantially pure compound, it may also be presented as a pharmaceutical formulation or preparation. The formulations to be used in the present invention, for both humans and animals, comprise any of the somatostatin agonists to be described below, together with one or more pharmaceutically acceptable carriers thereof, and
10 optionally other therapeutic ingredients.

 The carrier must be "acceptable" in the sense of being compatible with the active ingredient(s) of the formulation (e.g., capable of stabilizing peptides) and not deleterious to the subject to be treated. Desirably, the formulation should not include oxidizing agents or other substances with which peptides are known to be incompatible. For
15 example, somatostatin agonists in the cyclized form (e.g., internal cysteine disulfide bond) can be oxidized; thus, the presence of reducing agents as excipients could lead to an opening of the cysteine disulfide bridge. On the other hand, highly oxidative conditions can lead to the formation of cysteine sulfoxide and to the oxidation of tryptophane. Consequently, it is important to carefully select the excipient. pH is
20 another key factor, and it may be necessary to buffer the product under slightly acidic conditions (pH 5 to 6).

 The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient(s) into association with the carrier
25 which constitutes one or more accessory ingredients.

 In general, the formulations for tablets or powders are prepared by uniformly and intimately blending the active ingredient with finely divided solid carriers, and then, if necessary, as in the case of tablets, forming the product into the desired shape and size.

30 Formulations suitable for parenteral (e.g., intravenous) administration, on the other hand, conveniently comprise sterile aqueous solutions of the active ingredient(s). Preferably, the solutions are isotonic with the blood of the subject to be treated. Such formulations may be conveniently prepared by dissolving solid active ingredient(s) in water to produce an aqueous solution, and rendering said solution sterile. The
35 formulation may be presented in unit or multi-dose containers, for example, sealed ampoules or vials.

Formulations suitable for sustained release parenteral administrations (e.g., biodegradable polymer formulations such as polyesters containing lactic or glycolic acid residues) are also well known in the art. See, e.g., U.S. Patent Nos. 3,773,919 and 4,767,628 and PCT Publication No. WO 94/15587.

5 Methods and formulations for the treatment and/or coating of surgical stents, wherein said coating may comprise a pharmaceutically active compound, are also well known in the art. See, e.g., US Patents 6,214,115, 6,090,901 and 6,083,257, and International Patent Publication No.'s WO 01/01957 and WO 00/02599.

10 The somatostatin or somatostatin agonist may also be administered with another compound capable of lowering blood levels of triglycerides, cholesterol, or glycerol, such as fibrates (e.g., bezafibrate, gemfibrozil, and clofibrate), HMG-CoA reductase inhibitors (e.g., pravastatin, simvastatin, and fluvastatin, Atorvastatin, and Lovastatin), bile acid binding resins (e.g., cholestyramine and colestipol), nicotinic acid compounds (e.g., nicotinic acid and nicheritol), and fish oils. See Workshop Treatment of Hyperlipidemia
15 1996-2 (Lakemedelsverket, Uppsala, Sweden, 1996).

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments and from the claims.

DETAILED DESCRIPTION OF THE INVENTION

20 It is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent. The following specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

25 Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Also, all publications, patent applications, patents, and other references mentioned herein are incorporated by reference.

30 What is meant by a somatostatin type-1 receptor agonist (i.e., SSTR-1 agonist) is a compound which (1) has a high binding affinity (e.g., K_i of less than 1000 nM or preferably less than 100 nM or less than 10 nM) for SSTR-1 (e.g., as defined by the receptor binding assay described below) and (2) decreases the rate or extent of vascular proliferation, (e.g., as shown by the biological assay described below).

35 What is meant by a somatostatin type-1 receptor selective agonist is a somatostatin agonist which (1) has a higher binding affinity (i.e., K_i) for SSTR-1 than for either SSTR-2, SSTR-3, SSTR-4 or SSTR-5, (e.g., as defined by the receptor binding assay described below) and (2) decreases the rate or extent of vascular proliferation, (e.g., as shown by the biological assay described below).

In one embodiment, the somatostatin type-1 receptor selective agonist is also a SSTR-1 agonist.

Examples of somatostatin agonists are those covered by formulae or those specifically recited in the publications set forth below, all of which are hereby
5 incorporated by reference.

Van Binst, G. et al. Peptide Research 5:8 (1992);

Horvath, A. et al. Abstract, "Conformations of Somatostatin Analogs Having Antitumor Activity", 22nd European peptide Symposium, September 13-19, 1992, Interlaken, Switzerland;

10 Curtis et al., Am. J. Physiol. Heart. Circ. Physiol, 278:H1815 (2000);

Nicolaou et al., Design and synthesis of a peptidomimetic employing β -D-glucose for scaffolding, in Peptides, Rivier and Marshall, eds., ESCOM (1990),

Papageorgiou et al., "Design, synthesis, and binding affinity of a non peptide mimic of somatostatin" Bioorganic & Medicinal Chemistry Letters, vol. 2, pp. 135-140,
15 1992; and R. Hirschmann et al. "De novo design and synthesis of somatostatin non-peptide peptidomimetics utilizing beta-D-glucose as a novel scaffolding, J. Am. Chem. Soc., vol. 115, pp. 12550-12568, 1993.

PCT Application No. WO 91/09056 (1991);

EP Application No. 0 363 589 A2 (1990);

20 EP Application No. P5 164 EU (Inventor: G. Keri);

U.S. Patent No. 6,262,229

U.S. Patent No. 6,197,963

U.S. Patent No. 6,159,941

U.S. Patent No. 6,127,343

25 U.S. Patent No. 6,083,960

U.S. Patent No. 6,020,349

U.S. Patent No. 5,552,534

U.S. Patent No. 5,817,879

U.S. Patent No. 5,811,512

30 U.S. Patent No. 4,904,642 (1990);

U.S. Patent No. 4,871,717 (1989);

U.S. Patent No. 4,853,371 (1989);

U.S. Patent No. 4,725,577 (1988);

U.S. Patent No. 4,684,620 (1987);

35 U.S. Patent No. 4,650,787 (1987);

U.S. Patent No. 4,603,120 (1986);


U.S. Patent No. 4,585,755 (1986);

EP Application No. 0 203 031 A2 (1986);
U.S. Patent No. 4,522,813 (1985);
U.S. Patent No. 4,486,415 (1984);
U.S. Patent No. 4,485,101 (1984);
5 U.S. Patent No. 4,435,385 (1984);
U.S. Patent No. 4,395,403 (1983);
U.S. Patent No. 4,369,179 (1983);
U.S. Patent No. 4,360,516 (1982);
U.S. Patent No. 4,358,439 (1982);
10 U.S. Patent No. 4,328,214 (1982);
U.S. Patent No. 4,316,890 (1982);
U.S. Patent No. 4,310,518 (1982);
U.S. Patent No. 4,291,022 (1981);
U.S. Patent No. 4,238,481 (1980);
15 U.S. Patent No. 4,235,886 (1980);
U.S. Patent No. 4,224,199 (1980);
U.S. Patent No. 4,211,693 (1980);
U.S. Patent No. 4,190,648 (1980);
U.S. Patent No. 4,146,612 (1979);
20 U.S. Patent No. 4,133,782 (1979);
U.S. Patent No. 5,506,339 (1996);
U.S. Patent No. 4,261,885 (1981);
U.S. Patent No. 4,728,638 (1988);
U.S. Patent No. 4,282,143 (1981);
25 U.S. Patent No. 4,215,039 (1980);
U.S. Patent No. 4,209,426 (1980);
U.S. Patent No. 4,190,575 (1980);
EP Patent No. 0 389 180 (1990);
EP Application No. 0 505 680 (1982);
30 EP Application No. 0 083 305 (1982);
EP Application No. 0 030 920 (1980);
PCT Application No. WO 88/05052 (1988);
PCT Application No. WO 90/12811 (1990);
PCT Application No. WO 97/01579 (1997);
35 PCT Application No. WO 91/18016 (1991);
PCT Application No. WO 00/75186 (2000);
U.K. Application No. GB 2,095,261 (1981); and

French Application No. FR 2,522,655 (1983).


Examples of SSTR-1 selective somatostatin agonists include, but are not limited to, the following somatostatin analogs which are disclosed in the above-cited references:

- 5 H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂;
 H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂;
 H-Cys-Phe-Phe-D-Trp-Lys-Ser-Phe-Cys-NH₂;
 H-Cys-Phe-Tyr-D-Trp-Lys-Thr-Phe-Cys-NH₂;
 H-Cys-Phe-Tyr(I)-D-Trp-Lys-Thr-Phe-Cys-NH₂;



;
 and

Caeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂, in which the structure for "Caeg" is



Note that for all somatostatin agonists described herein, each amino acid residue represents the structure of -NH-C(R)H-CO-, in which R is the side chain (e.g., CH₃ for Ala). Lines between amino acid residues represent peptide bonds which join the amino acids. Also, where the amino acid residue is optically active, it is the L-form configuration that is intended unless D-form is expressly designated. A disulfide bond (e.g., a disulfide bridge) exists between the two free thiols of the Cys residues; however, it is not shown.

Synthesis of somatostatin agonists

The methods for synthesizing somatostatin agonists is well documented and are within the ability of a person of ordinary skill in the art. For example, synthesis of H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂, described above, can be achieved by following the protocol set forth in Example I of European Patent Application 0 395 417 A1. The synthesis of somatostatin agonists with a substituted N-terminus can be achieved, for example, by following the protocol set forth in WO 88/02756, European Patent Application No. 0 329 295, and PCT Publication No. WO 94/04752.

Somatostatin Receptor Binding Assays

The human SSTR-1, SSTR-2, SSTR-3, SSTR-4, and SSTR-5 cDNA clones have been described (SSTR-1 and SSTR-2 in Yamada, Y., et al., Proc. Natl. Acad. Sci. USA., 89:251-255 (1992); SSTR-3 in Yamada, et al., Mol. Endocrinol. 6:2136-2142 (1993); and SSTR-4 and SSTR-5 in Yamada, et al., Biochem. Biophys. Res. Commun. 195:844-852 (1993)) and are also available from American Type Culture Collection (ATCC, Rockville, MD) (ATCC Nos. 79044 (SSTR-1), 79046 (SSTR-2), and 79048 (SSTR-3)). Based on the restriction endonuclease maps, the entire coding region of each SSTR cDNA may be excised by suitable restriction endonuclease digestion (Maniatis, T., et al., *Molecular Cloning - A Laboratory Manual*, CSHL, 1982). Restriction endonucleases are available from New England Biolabs (Beverly, MA). This cDNA fragment was inserted into the mammalian expression vector, pCMV (Russell, D., et al., J. Biol. Chem., 264:8222-8229 (1989)), using standard molecular biology techniques (see e.g., Maniatis, T., et al., *Molecular Cloning, -A Laboratory Manual*, Cold Spring Harbor Laboratory, 1982) to produce the expression plasmid, pCMV-human SSTR-1 through pCMV-human SSTR-5. Other mammalian expression vectors include pcDNA1/Amp (Invitrogen, Sandlesy, CA). The expression plasmids were introduced into the suitable bacterial host, E. Coli HB101 (Stratagene, La Jolla, CA) and plasmid DNAs, for transfection, were prepared on Cesium Chloride gradients.

CHO-K1 (ovary, Chinese hamster) cells were obtained from ATCC (ATCC No. CCL 61). The cells were grown and maintained in Ham's F12 media (Gibco BRL, Grand Island, NY) supplemented with 10% fetal bovine serum under standard tissue culture conditions. For transfection, the cells were seeded at a density 1×10^6 /60-cm plate (Baxter Scientific Products, McGraw Park, IL.). DNA mediated transfection was carried out using the calcium phosphate co-precipitation method (Ausubel, F.M., et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, 1987). The plasmid pRSV-neo (ATCC; ATCC No. 37198) was included as a selectable marker at 1/10 the concentration of the expression plasmid. CHO-K1 clonal cell lines that have stably inherited the transfected DNA were selected for growth in Ham's F12 media containing 10% fetal bovine serum and 0.5mg/ml of G418 (Sigma). The cells were ring-cloned and expanded in the same media for analysis.

Expression of the human SSTR-1 through SSTR-5 receptors in the CHO-K1 cells were detected by Northern blot analysis of total RNA prepared from the cells (Sambrook, J.E., et al., *Molecular Cloning - A Laboratory Manual*, Ed. 2., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989) and by receptor binding using (125 I-Tyr¹¹)somatostatin-14 as a ligand. Transfected cell lines expressing the human SSTR receptors were clonally expanded in culture and used in the following SSTR binding

protocol.

Crude membranes were prepared by homogenization of the transfected cells in 20 ml of ice-cold 50 mM Tris-HCl with a POLYTRON homogenizer (setting 6, 15 sec). Buffer was added to obtain a final volume of 40 ml, and the homogenate was centrifuged in a
5 Sorval SS-34 rotor at 39,000 g for 10 min at 0-4°C. The resulting supernatant was decanted and discarded. The pellet was rehomogenized in ice-cold buffer, diluted, and centrifuged as before. The final pellet was resuspended in the 10 mM Tris HCl and held on ice for the receptor binding assay.

Aliquots of the membrane preparation were incubated for 30 min at 30°C with
10 0.05 nM (¹²⁵I-Tyr¹¹)somatostatin-14 (2000 Ci/mmol; Amersham Corp., Arlington Heights, IL) in 50 mM HEPES (pH 7.4) containing a test somatostatin agonist of various concentrations (e.g., 10⁻¹¹ to 10⁻⁶), 10 mg/ml bovine serum albumin (fraction V) (Sigma Chemical Co., St. Louis, MO), MgCl₂ (5 mM), Trasylol (200 KIU ml), bacitracin (0.02 mg/ml), and phenylmethylsulphonyl fluoride (0.02 mg/ml). The final assay volume was
15 0.3 ml. The incubations were terminated by rapid filtration through GF/C filters (pre-soaked in 0.3% polyethylenimine for 30 min) using a Brandel filtration manifold. Each tube and filter were then washed three times with 5 ml aliquots of ice-cold buffer. Specific binding was defined as the total (¹²⁵I-Tyr¹¹)SRIF-14 bound minus that bound in the presence of 1000 nM. The K_i values for somatostatin agonists are calculated by
20 using the following formula: $K_i = IC_{50} / (1 + (LC/LEC))$ where IC₅₀ is the concentration of test somatostatin agonist required to inhibit 50 percent of the specific binding of the radioligand (¹²⁵I-Tyr¹¹)somatostatin-14, LC is the concentration of the radioligand (0.05 nM), and LEC is the equilibrium dissociation constant of the radioligand (0.16 nM).

Inhibition of Proliferation and Capillary Tube Formation

25 In order to investigate the antiproliferative effects on endothelial cells and the inhibition of proliferation and capillary tube formation of endothelial cells caused by somatostatin analogues, two different human *in vitro* models have been preformed. These models permitted study of the effects of somatostatin analogues on a bi-dimensional endothelial cell layer and on three-dimensional endothelial cell growth in
30 an extracellular matrix that mimic the capillary development *in vivo*. Furthermore, these assays allowed a very long period of treatment (72 hours and 28 days, respectively) on human tissues, resembling the possible chronic clinical approach to antiangiogenic therapy.

Material and methods**Materials**

Recombinant human epidermal growth factor (EGF) and recombinant human vascular endothelial growth factor (VEGF) were from PeproTechEC LTD (London, UK).

- 5 EGF and VEGF, according with the information of data sheet, were reconstituted in sterile distilled water at a concentration of 100 µg/mL.

- Cell culture medium 199 and medium 199 without phenol red were purchased from Gibco BRL (Paisley, UK). Type A gelatin from porcine skin supplements and all other chemicals not listed in this section were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Plastics for cell culture were supplied by Costar (Cambridge, MA, USA).

- Somatostatin-14, BIM-23014C, BIM-23120C, BIM-23190C, BIM-23197C, BIM-23206C, BIM-23268C, BIM-23745C and BIM-23926C (each from Biomeasure, Incorporated, Milford, MA, USA) were dissolved in a stock solution of 0.01N acetic acid containing 0.1% fatty acid-free bovine serum albumin (BSA) and stored at -80°C.

The relative affinities of the foregoing compounds for the various somatostatin receptors may be summarized as follows:

Compound	SSTR-1	SSTR-2	SSTR-3	SSTR-4	SSTR-5
20 Somatostatin-14	v. high	v. high	v. high	v. high	v. high
BIM-23014C	v. low	v. high	low	v. low	high
BIM-23120C	v. low	v. high	v. low	v. low	low
BIM-23190C	v. low	v. high	low	v. low	high
25 BIM-23197C	v. low	v. high	mod-high	v. low	high
BIM-23206C	v. low	low	v. low	v. low	v. high
BIM-23268C	high	mod-high	high	mod-high	v. high
30 BIM-23745C	mod	v. low	v. low	v. low	v. low
BIM-23926C	v. high	v. low	v. low	v. low	v. low

SU5416, 3-((2,4-dimethylpyrrol-5-yl) methylidenyl)-2-indolinone, was a gift from Sugen Inc. (San Francisco, CA, USA).

Cell culture conditions

- 35 The immortalized human microvascular endothelial cell line HMEC-1, characterized by Ades, *et al.*, (Journal of Investigative Dermatology 1992, 99: 683-690), was maintained in Medium 199, supplemented with 10% heat-inactivated fetal bovine

serum (FBS), penicillin (50 IU/mL) and hydrocortisone (100 µg/mL). Cells were routinely grown in 75 cm² gelatin-coated tissue culture flasks and kept in a humidified atmosphere of 5% CO₂ at 37°C. Cells were harvested with a solution of 0.25% trypsin-0.03% EDTA when they were in long phase of growth and maintained at the above-described culture conditions for all experiments.

Cytotoxicity assay

In vitro chemosensitivity testing was performed on single-cell suspensions of HEMC-1 cells (5x10³ cells/well) plated in 96-well gelatin-coated sterile plastic plates and allowed to attach overnight. The treatment protocol (figure 1) was designed so that after 24 hours, 10⁻¹⁰-10⁻⁶ M somatostatin-14, somatostatin analogues, SU5416 10⁻⁶ M + VEGF 10 ng/mL as positive control or vehicle were added, and plates were incubated for 72 hours (for more details, see figure 1). The treatments were scheduled so that each peptide was represented by at least nine wells. At the end of the experiment, cells were rinsed with phosphate buffered saline (PBS), harvested with trypsin/EDTA, and counted with an hemocytometer. Results are expressed as the percentage of cell proliferation versus controls and are the mean of three separate experiments ± S.E repeated thrice.

In vitro cultures of human placental vessels

The use of the experimental model of human placental explants, detailed below, received authorization by the Ethics committee of Pisa University Hospital (Protocol n. 005567).

The experimental procedure described by Brown et al., (Laboratory Investigation 1996, 75: 539-555) was followed and modified in the present study. Immediately after the spontaneous delivery, the placenta was collected in sterile conditions and superficial blood vessels, approximately 1 to 1.5 mm in diameter and from 1 to 5 cm in length, were excised. Vessel explants were placed in phosphate buffered saline (PBS) solution containing 2.5 mg/mL of amphotericin B and 50 µg/mL of gentamycin and were cut into approximately 1-mm fragments. The cultures were performed in 24-well culture plates; 0.5 mL/well of a solution of fibrinogen 3 mg/mL in Medium 199 without phenol red was added to each well followed by the quick addition of 15 µL of thrombin (50 NIH U/mL in 0.15 M NaCl). The vessel explants were rapidly placed in the center of the wells after clot formation and covered by 0.5 mL/well of the fibrinogen solution with the addition of further 15 µL of thrombin for suspending all of them at the same level between the two clots. After gel formation, 1 mL/well of medium 199 without phenol red supplemented with 10% of heat-inactivated FBS, 0.1% ε-aminocaproic acid, L-glutamine (2 mM), and antibiotics (streptomycin 50 µg/mL, penicillin 50 IU/mL and amphotericin B 2.5 mg/mL) were added. Vessels were cultured and treated every two days with 10⁻¹⁰-10⁻⁶ M

somatostatin-14, somatostatin analogues, and SU5416 10^{-6} M or vehicle at 37°C 95% air/5% CO₂ in a humidified environment for 28 days (figure 2). Vessels explants were photographed on day 28 with a phase-contrast Leitz MD IL microscope (Leica, Heerbrugg, Switzerland) and were subjected to image analysis.

5 *Image Analysis*

The image analysis procedures described by Bocci *et al.*, (Cancer Chemotherapy Pharmacology 1999, 43: 205-212), were adopted for the present study. Briefly, photographs obtained from the placental fragment assay were digitized in a 512 x 512-pixel matrix, using a color video camera TK-1280E (JVC, Tokyo, Japan) and a
10 microcomputer processor. Digitized pictures were visualized on highresolution color display. The true color image analysis software package KS 300 v.1.2 (Kontron Elektronik GmbH, Eching, Germany) was run for interactive manipulation, quantification of the images and data collection. Geometric calibrations were set with a sample of
15 the image that was in the range of 0-255, where 0 was black (presence of vascular sprouts) and 255 was white (absence of vascular sprouts). In the fibrin culture of placental vessel explants the mean gray level of the sprouting area was measured and the sprouting index (SI) was defined as:

20
$$\text{Sprouting index} = ((\text{sprouting area}/\text{mean gray level of sprouting area})/\text{perimeter of explant}) \times 100.$$

Results are expressed as the percentage of sprouting index +/- S.E. *versus* controls.

Results

Cytotoxicity assay

All results are shown in Table I. All somatostatin analogues at the studies
25 concentrations revealed antiproliferative activity on immortalized human microvascular endothelial HMEC-1 cells, with a maximum effect at 10^{-7} – 10^{-8} M. The positive control of SU5416 10^{-6} M, a specific VEGF-receptor inhibitor, resulted in a cell growth block of 56.2%.

In vitro cultures of human placental vessels

30 The explant sprouting within the fibrin matrix was characterized by numerous microvessels around the placental fragment. Vascular cells organized radially to form microvessels that underwent continuous remodeling (figure 3). The maximal growth of the three-dimensional microvascular network occurred during the third-fourth week and reached the plateau at 27 days after explant. Histologically, in the fibrin gel a subtle
35 framework of endothelial cells, that was immunoreactive for von Willebrand factor (figure 4), was observed (figure 5). In most of the cases the microvessels showed an initial

lumen; in other the lumen was absent and only endothelial cells were observed (figure 5).

The microscopic picture of the placental explants on the first day of the culture is shown in figure 6 A (bar, 2 mm); the appearance of the outgrowth of endothelial cells from the placental cultured vessel fragment was observed approximately on the sixth day of culture ($SI=0.055\pm0.004$ (mm/mean gray) x 100; figure 6 D).

The experimental data on the activity of somatostatin analogues on capillaries sprouts were summarized in Table II. Cultures treated with SMS analogues and SU5416 were shown in figures 7-16 where the maximum effects were shown. BIM-23926C and BIM-23745C revealed a potent inhibitory property in a long-term treatment; they resulted, respectively, in $17.18\pm11.8\%$ at 10^{-7} M and $42.84\pm5.6\%$ at 10^{-8} M of SI as compared to untreated controls. SU5416, the positive control, resulted in $32.92\pm9.7\%$ of SI at a concentration of 10^{-8} M.

OTHER EMBODIMENTS

The foregoing description has been limited to specific embodiments of this invention. It will be apparent, however, that variations and modifications may be made to the invention, with the attainment of some or all of the advantages of the invention. Such embodiments are also within the scope of the following claims.

What is claimed is:

CLAIMS


1. A method of treating vascular proliferation in a patient, said method
 5 comprising administering a therapeutically effective amount of a somatostatin type-1
 receptor agonist to said patient.

2. The method of claim 1, wherein said somatostatin type-1 receptor
 agonist has a K_i of less than 5 nM for the somatostatin type-1 receptor.
 10

3. The method of claim 1, wherein said wherein said somatostatin type-1
 receptor agonist is a somatostatin type-1 receptor selective agonist.

4. The method of claim 3, wherein said somatostatin type-1 receptor
 15 selective agonist has a K_i for the type-1 somatostatin receptor that is at least 10 times
 less than its K_i for each of the somatostatin type-2, type-3, type-4, and type-5 receptors.

5. The method according to claim 1 wherein the somatostatin type-1
 receptor agonist is :
 20 H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂;
 H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂;
 H-Cys-Phe-Phe-D-Trp-Lys-Ser-Phe-Cys-NH₂;
 H-Cys-Phe-Tyr-D-Trp-Lys-Thr-Phe-Cys-NH₂;
 H-Cys-Phe-Tyr(I)-D-Trp-Lys-Thr-Phe-Cys-NH₂;

HO(CH₂)₂N  N-(CH₂)-CO-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂
 25 ;

Caeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂; or
 3-((2,4-dimethylpyrrol-5-yl) methylidenyl)-2-indolinone;
 or a pharmaceutically acceptable salt thereof.

6. The method of claim 5, said method comprising administering a
 therapeutically effective amount of a compound according to the formula:

Caeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 or a pharmaceutically acceptable salt thereof.
 35

7. The method of claim 1, wherein said vascular proliferation comprises angiogenesis, restenosis, endothelial cell proliferation, smooth muscle proliferation, or new blood vessel sprouting.

5 8. The method of claim 1, wherein said vascular proliferation occurs in a disease or condition comprising an autoimmune disease, arthritis, scleroderma, cancerous tumors, corneal graft neovascularization, diabetic retinopathy, hemangioma, hypertrophic scarring, or psoriasis.

10 9. The method of claim 1, wherein said vascular proliferation is incident to or associated with angioplasty or an AV shunt.

10. The method of claim 1, wherein said vascular proliferation occurs in a disease or condition comprising warts, granulomas, Kaposi's sarcoma, or allergic oedema.

11. The method of claim 1, wherein said vascular proliferation occurs in a disease or condition comprising endometriosis, dysfunctional uterine bleeding, or follicular cysts.

20 12. The method of claim 1, wherein said vascular proliferation occurs in a disease or condition comprising retinopathy of prematurity, choroidopathy, macular degeneration, or age-related macular degeneration.

25 13. The method of claim 1, wherein said vascular proliferation occurs in a disease or condition comprising solid tumor, tumor metastasis, benign tumor, acoustic neuromas, neurofibromas, trachomas, leukemia, pyogenic granuloma, myocardial angiogenesis, plaque neovascularization, atherosclerosis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, corneal graft rejection, Osler-Webber Syndrome, rubeosis, neovascular glaucoma, retrolental
30 fibroplasia, diabetic retinopathy, diabetic neovascularization, fracture, vasculogenesis, hematopoiesis, ovulation, menstruation, placentation, cat scratch disease (Rochele minalia quintosa), peptic ulcer, Helicobacter pylori associated ulcer, psoriasis, telangiectasia psoriasis, rheumatoid arthritis, Crohn's disease, intestinal adhesions,
35 scarring, hypertrophic scars, keloids, telangiectasia, hemophiliac joints, angiofibroma, or wound granulation.

14. The method of claim 1, wherein said somatostatin type-1 receptor agonist is disposed upon or within a vascular stent.

15. The method of claim 14, wherein said somatostatin type-1 receptor agonist is
5 provided as a component of a slow release formulation.

16. The method of claim 14, wherein said somatostatin type-1 receptor agonist is provided as a component of a polymeric composition.




Figure 1. Treatment protocol of HEMC-1 cells




Figure 2. Treatment protocol of placental vessel explants




Figure 3



Figure 5






Figure 4

Figure 6




A



B



C



D




Figure 7. SMS-14 10^{-7} M




Figure 10. BIM23190C 10^{-9} M



Figure 9. BIM23745C 10^{-8} M

Figure 8. BIM23014C 10^{-6} M




Figure 12. BIM23206C 10^{-8} M



Figure 14. BIM23120C 10^{-8} M




Figure 11. BIM23268C 10^{-7} M



Figure 13. BIM23197C 10^{-7} M



Figure 15. BIM23928C 10^{-7} M



Figure 16. SU5416 10^{-6} M

Table 1 Proliferation of HEMC-1 human endothelial cells *in vitro*

Concentration	BIM-23120C	BIM-23197C	BIM-23206C	BIM-23190C	BIM-23268C	BIM-23745C	BIM-23014C	BIM-23926C	SMS-14
0 M	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100
10 ⁻⁶ M	103.97 ± 2.07	87.61 ± 3.6	122.58 ± 7.3	113.55 ± 6.5	118.06 ± 8.2	40.0 ± 2.0	80.0 ± 3.2	56.78 ± 4.9	73.3 ± 4.1
10 ⁻⁷ M	74.27 ± 4.5	78.25 ± 2.5	81.29 ± 3.2	92.26 ± 3.1	87.74 ± 3.7	46.55 ± 2.3	51.2 ± 4.7	55.33 ± 3.3	71.25 ± 2.1
10 ⁻⁸ M	88.01 ± 3.7	73.39 ± 2.1	68.38 ± 3.9	60.0 ± 2.2	111.61 ± 2.5	49.7 ± 1.9	65.05 ± 3.1	65.54 ± 3.5	69.7 ± 2.5
10 ⁻⁹ M	102.39 ± 5.2	83.96 ± 3.2	77.42 ± 4.0	87.74 ± 2.5	94.19 ± 4.2	56.79 ± 2.7	78.0 ± 2.7	79.95 ± 4.6	77.85 ± 3.3

Treatment was delivered on day 1, 24 hours after plating and repeated 48 hours later; cell count was performed after additional 24 hours. SMS-14 was added on each day. SU5416 was added at 10⁻⁶ M to culture media of separate wells and resulted in 43.85 ± 3.02 % survival as compared to untreated controls.

Table II Proliferation of endothelial cells from blood vessels explanted from human placenta

Concentration	BIM-23120C	BIM-23197C	BIM-23206C	BIM-23190C	BIM-23268C	BIM-23745C	BIM-23014C	BIM-23926C	SMS-14
0 M	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100	% of controls 100
10 ⁻⁶ M	106.6 ± 10.2	98.99 ± 8.7	93.08 ± 7.8	107.98 ± 8.5	98.14 ± 10.7	70.75 ± 7.5	73.66 ± 8.7	38.39 ± 10.6	73.25 ± 7.5
10 ⁻⁷ M	94.78 ± 8.4	62.72 ± 7.9	83.77 ± 10.2	102.39 ± 7.3	81.12 ± 11.6	62.99 ± 8.2	79.01 ± 8.2	17.18 ± 11.8	44.03 ± 5.2
10 ⁻⁸ M	88.18 ± 11.3	71.54 ± 8.2	72.07 ± 8.5	95.12 ± 10.7	84.31 ± 12.1	42.84 ± 5.6	80.24 ± 10.1	41.54 ± 7.9	68.72 ± 5.4
10 ⁻⁹ M	78.76 ± 7.6	114.23 ± 7.6	97.87 ± 10.9	91.75 ± 10.9	86.97 ± 8.4	63.41 ± 6.8	90.33 ± 11.3	55.94 ± 13.6	84.77 ± 7.1
10 ⁻¹⁰ M	86.57 ± 9.8	126.65 ± 11.2	105.32 ± 11.3	103.98 ± 12	76.33 ± 7.6	82.43 ± 8.5	90.94 ± 12.6	62.32 ± 12	99.18 ± 2

Treatments were delivered every after day over a 28-day period. SU5416 was added every two days at 10⁻⁶ M culture media of separate explants and resulted in 32.92 ± 9.7 % of SI as compared to untreated controls

Figure 17

This Page Blank (uspto)

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 02/01125

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/31 A61K31/404 A61P43/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data, EMBASE, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>BARRIE R ET AL: "INHIBITION OF ANGIOGENESIS BY SOMATOSTATIN AND SOMATOSTATIN-LIKE COMPOUNDS IS STRUCTURALLY DEPENDENT" JOURNAL OF SURGICAL RESEARCH, ACADEMIC PRESS INC., SAN DIEGO, CA, US, vol. 4, no. 55, 1993, pages 446-450, XP001063285 ISSN: 0022-4804 see discussion figures 1,2</p> <p style="text-align: center;">-- -/-</p>	1-4, 7, 8, 12, 13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

10 March 2003

Date of mailing of the international search report

14/03/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/US 02/01125

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MCCOMBE M ET AL: "EFFECT OF A LONG-ACTING SOMATOSTATIN ANALOGUE (BIM23014) ON PROLIFERATIVE DIABETIC RETINOPATHY: A PILOT STUDY" EYE, ROYAL COLLEGE OF OPHTHALMOLOGISTS, LONDON, GB, vol. 5, no. 5, 1991, pages 569-575, XP001031225 see discussion abstract	1-4,7,8, 12,13,15
T	AAVIK EINARI ET AL: "Elimination of vascular fibrointimal hyperplasia by somatostatin receptor 1,4-selective agonist." THE FASEB JOURNAL: OFFICIAL PUBLICATION OF THE FEDERATION OF AMERICAN SOCIETIES FOR EXPERIMENTAL BIOLOGY. UNITED STATES MAY 2002, vol. 16, no. 7, May 2002 (2002-05), pages 724-726, XP008014600 ISSN: 1530-6860 see conclusions and significance abstract	1-4,7,9
X	WO 99 49884 A (PATEL YOGESH C ;HAEYRY PEKKA (FI); JUVANTIA PHARMA LTD OY (FI)) 7 October 1999 (1999-10-07) page 5; claims 1-4,7; table 1 page 6, line 20-30	1-4,7-9, 12-14
X	WO 00 12111 A (UNIV BRITISH COLUMBIA ;BUCHAN ALISON (CA); HSIANG YORK (CA); LEVY) 9 March 2000 (2000-03-09) abstract; claims 1,4,6,8 page 21, line 11-20	1-4,7,8, 13,15,16
X	WO 99 56769 A (SOD CONSEILS RECH APPLIC ;KANEKO HIROSHI (JP); MITSUMA TERUNORI (J) 11 November 1999 (1999-11-11) page 29, line 15.25; claims 13,14,17,18,31; examples 108-110,118	1-5,7,8, 13,15,16
X	WO 99 65508 A (GELLERMAN GARY ;HORNIK VERED (IL); PEPTOR LTD (IL); AFARGAN MICHEL) 23 December 1999 (1999-12-23) page 59; claims 35,35	1-5,7,8, 12,13
A	WO 96 07398 A (CHERIF CHEIKH ROLAND ;SOD CONSEILS RECH APPLIC (FR)) 14 March 1996 (1996-03-14) claims 1,2	1-16

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/01125

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-16 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: partially 1-16
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Claims Nos.: partially 1-16

Present claims 1-16 relate to compounds/compositions defined by reference to desirable characteristic or properties, namely: "a somatostatin type 1 receptor agonist" (claim 1); "a somatostatin type 1 receptor selective agonist" (claim 3); "a slow release formulation " (claim 15); "a polymeric composition" (claim 16).

The claims cover all compounds/compositions having these characteristics or properties, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds/methods.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved: "a somatostatin type 1 receptor agonist" (claim 1); "a somatostatin type 1 receptor selective agonist" (claim 3); "a slow release formulation " (claim 15).

Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

Furthermore claims 2,4 relate to a compound defined (inter alia) by reference to the following parameter: "has a K_i of less than..." The use of this parameter in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT.

It is impossible to compare the parameter the applicant has chosen to employ with what is set out in the prior art.

Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds specifically used in the examples (page 12) and those mentioned in claims 5, 6 in relation to their use for the treatment of diseases as claimed in claims 7-13.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: Application No

PCT/US 02/01125

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9949884	A	07-10-1999	US 6124256 A CA 2323748 A1 EP 1064017 A1 WO 9949884 A1 JP 2002509894 T	26-09-2000 07-10-1999 03-01-2001 07-10-1999 02-04-2002
WO 0012111	A	09-03-2000	AU 5499799 A CA 2340588 A1 WO 0012111 A2 CN 1320042 T EP 1107780 A2 JP 2002523465 T NO 20011025 A US 2002137676 A1	21-03-2000 09-03-2000 09-03-2000 31-10-2001 20-06-2001 30-07-2002 30-03-2001 26-09-2002
WO 9956769	A	11-11-1999	US 5968903 A AU 3975499 A EP 1075273 A2 JP 2002513769 T NO 20005588 A WO 9956769 A2	19-10-1999 23-11-1999 14-02-2001 14-05-2002 05-01-2001 11-11-1999
WO 9965508	A	23-12-1999	US 6051554 A US 6355613 B1 AU 747515 B2 AU 4288499 A CA 2335488 A1 EP 1085896 A1 WO 9965508 A1 JP 2002518339 T US 2002052315 A1	18-04-2000 12-03-2002 16-05-2002 05-01-2000 23-12-1999 28-03-2001 23-12-1999 25-06-2002 02-05-2002
WO 9607398	A	14-03-1996	US 5595760 A US 5837276 A US 5582591 A AT 210424 T AU 713066 B2 AU 3398595 A BR 9508687 A CA 2198917 A1 CN 1165479 A DE 69524609 D1 DE 69524609 T2 DK 779805 T3 EP 1157703 A2 EP 0779805 A2 ES 2169762 T3 WO 9607398 A2 JP 10508294 T PT 779805 T AT 188615 T AU 3398295 A DE 69514530 D1 DE 69514530 T2 DK 778785 T3 EP 0778785 A2 EP 0953362 A2 ES 2141956 T3	21-01-1997 17-11-1998 10-12-1996 15-12-2001 25-11-1999 27-03-1996 02-06-1998 14-03-1996 19-11-1997 24-01-2002 21-11-2002 25-02-2002 28-11-2001 25-06-1997 16-07-2002 14-03-1996 18-08-1998 29-04-2002 15-01-2000 27-03-1996 17-02-2000 27-07-2000 29-05-2000 18-06-1997 03-11-1999 01-04-2000

INTERNATIONAL SEARCH REPORT

tion on patent family members

Interns Application No
PCT/US 02/01125

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9607398 A		GR 3032947 T3	31-07-2000
		WO 9607440 A2	14-03-1996
		PT 778785 T	28-04-2000
		US 6142972 A	07-11-2000
		US 5660846 A	26-08-1997
		US 6306420 B1	23-10-2001
		ZA 9507355 A	25-06-1996
		AT 230977 T	15-02-2003
		AU 712916 B2	18-11-1999
		AU 3398195 A	27-03-1996
		BR 9508688 A	28-10-1997
		CA 2198916 A1	14-03-1996
		CN 1165477 A	19-11-1997
		DE 69529415 D1	20-02-2003
		EP 1270032 A1	02-01-2003
		EP 0778767 A2	18-06-1997
		WO 9607397 A2	14-03-1996
		JP 10508293 T	18-08-1998
		US 5616123 A	01-04-1997
		ZA 9507356 A	02-04-1996